

Doped manganites as cathodes for sodium-ion batteries: a self-consistent DFT+U study

Domenico CORONA – *University of Rome Tor Vergata*

Sodium-ion batteries share similarities with the well-established lithium-ion batteries currently in widespread use; in fact, analogously to lithium-ion positive electrodes, sodium-ion cathodes store sodium through an intercalation mechanism. Cathodes based on P2-NaxTMO₂ crystals, i.e., sodium based layered transition-metal oxides (LTMOs), have attracted growing interest owing to their high potential and capacity. However, the geometric structure of LTMOs may undergo a Jahn-Teller distortion during charge/discharge cycles. Doping has proven effective in stabilizing the P2 geometric structure improving the cycling stability by limiting structural transitions as well as delaying them at higher voltages. Ab-initio calculations within the Quantum ESPRESSO suite are employed to determine the optimal dopants, the total energy of the system, and estimate the sodium-ion intercalation potential. The structural and electronic properties of doped LTMOs are investigated at a GGA+U level of theory, where the on-site Hubbard U parameters are evaluated self-consistently using linear-response theory, so to realistically include the screening from the environment which is crucial for quantitatively predictive calculations on novel materials.

Acknowledgments: This work has been financed by the Research Fund for the Italian Electrical System under the Three Year Research Plan 2022-2024 (DM MITE n. 337, 15.09.2022), in compliance with the Decree of April 16th, 2018”.